



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

RECENT MATHEMATICAL PUBLICATIONS.

COMMUNICATED BY G. W. HILL.

THE Uranian and Neptunian Systems, investigated with the 26-inch equatorial of the U. S. Naval Observatory. By Prof. S. Newcomb. (Appendix I, Wash. Obs. for 1873.) 4to. Washington, 1875. 74 pp.

Many of our text-books attribute 8 satellites to Uranus; it is however now conclusively established that but 4 have been seen. So far as observation shows, the orbits of all the satellites of these two planets are circular, and those of Uranus lie in one plane. As this valuable paper may not be accessible to many of the readers of the ANALYST, we append the final elements arrived at. T being the fraction of a century after 1850, the position of the common plane of the orbits of the satellites of Uranus with reference to the ecliptic is given by the following inclination and longitude of ascending node

$$i = 97^{\circ}.85 - 0^{\circ}.013 T,$$

$$\theta = 165^{\circ}.48 + 1^{\circ}.40 T.$$

And a denoting the angle under which the radius vector of the satellite is seen at a distance whose logarithm is 1.28310, P the sidereal period in mean solar days, and u the argument of the Uranocentric latitude of the satellite at the epoch 1872, Jan. 0. 0, Washington mean time, we have

	a .	P .	u .
Ariel	13''.78	2 ^d .520383	15 ^o .82
Umbriel	19 .20	4 .144180	130 .56
Titania	31 .48	8 .705897	224 .00
Oberon	42 .10	13 .463269	148 .97

The mass of Uranus which corresponds to these elements is $\frac{1}{32350}$.

In like manner the elements of the satellite of Neptune, a being for the distance whose logarithm is 1.47814, and u for the epoch 1874, Jan. 0. 0, Washington mean time, are

$$i = 145^{\circ}.12, \quad \theta = 184^{\circ}.50 + 1^{\circ}.40 T,$$

$$a = 16''.275, \quad P = 5^d.876900, \quad u = 101^{\circ}.07.$$

The corresponding mass of Neptune is $\frac{1}{19380}$. The paper contains brief tables founded on these elements.

Lessons on Rigid Dynamics, by Rev. G. Pirie. Macmillan. New York, 1875. 8vo. \$2.25.

Oeuvres de Lagrange, publiques par les soins de J. A. Serret, sous les auspices de S. Exc. le Ministre de l' Instruction Publique. Tomes V-VI. Paris, 1873-4. 4to. 30 fr. per vol.

Theorie des Fonctions Elliptiques, par MM. Briot et Bouquet. Deuxieme Edition. Gauthier-Villars. Paris, 1875. 4to. 30 fr.

Abhandlungen von F. W. Bessel. Herausgegeben von R. Engelmann. In 3 Banden. Band I. Leipzig, 1875. 4to. 376 pp.